

## **School Vaccination Medical Exemption Form**

The School Vaccination Medical Exemption Form is the official Division of Public Health (DPH) document to be completed by a currently licensed physician, advanced practice nurse, nurse practitioner, or physician's assistant to exempt a child from childcare or school immunization requirements. The clinician certifies that due to the child's health or medical condition, the child may be adversely affected on a temporary or permanent basis by one or more of the required vaccines. The signed medical exemption statement verifying true contraindications/precautions is submitted to and accepted by Delaware schools, child care programs, and other agencies that require proof of immunization. The signed form does not require further approval from the Delaware Division of Public Health.

For exemption of medical conditions not listed on this form, the clinician must submit the Supplemental School Vaccine Medical Exemption Form to DPH for approval.

A *contraindication* is a condition in a recipient that increases the risk for a serious adverse reaction. A vaccine will not be administered when a contraindication exists.

A **precaution** is a condition in a recipient that might increase the risk for a serious adverse reaction or compromise the ability of the vaccine to produce immunity. Under normal conditions, vaccinations should be deferred when a precaution is present. Indicate if an exemption is permanent or temporary.

Vaccine medical contraindications are determined by the Advisory Committee on Immunization Practices (ACIP).

Please return the form to:	
School:	
Address:	
Fax Number:	



Please indicate whether the exemption is:   permanent	or	☐ <u>temporary</u>	
For <i>temporary</i> exemption, list the date the exemption ends	:	/ /	

Vaccine	Check if Applicable		Contraindications/Precautions
DTaP	Permanent	Temporary	Contraindications
			Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
			Encephalopathy within seven days after receipt of previous dose of DTP or DTaP
			Precautions
			<ul> <li>Progressive neurologic disorder, including infantile, spasms, uncontrolled epilepsy, progressive encephalopathy: defer DTaP until neurologic status clarified and stabilized</li> </ul>
			● Fever greater than 104.9°F (40.5°C) within 48 hours after vaccination of previous dose of DTP/DTaP
			Guillain-Barre syndrome within 6 weeks after a previous dose of tetanus toxoid-containing vaccine
			Hypotonic-hyporesponsive episodes within 48 hours after vaccination of previous dose of DTP or DTaP
			Seizure within 72 hours after vaccination of previous dose of DTP or DTaP
			<ul> <li>Persistent, inconsolable crying lasting three hours or more 48 hours after receiving a previous dose of DTP or DTaP</li> </ul>
			Moderate or severe acute illness with or without fever
			<ul> <li>History of arthus-type hypersensitivity reactions after a previous dose of tetanus or diphtheria toxoid-containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid containing vaccine</li> </ul>
Tdap			Contraindications
			Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
			Encephalopathy within seven days after receipt of previous dose of DTP or DTaP
			Precautions
			<ul> <li>Progressive neurologic disorder, uncontrolled seizures or progressive encephalopathy; defer vaccination until treatment regimen is established and condition stabilized.</li> </ul>
			Moderate or severe acute illness with or without fever
			Guillain-Barre syndrome within 6 weeks after a previous dose of tetanus toxoid-containing vaccine
			<ul> <li>History of arthus-type hypersensitivity reactions after a previous dose of tetanus or diphtheria toxoid-containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid containing vaccine</li> </ul>
DT/Td			Contraindications
			Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
			Precautions
			Guillain-Barre syndrome within 6 weeks after a previous dose of tetanus toxoid-containing vaccine
			Moderate or severe acute illness with or without fever
			<ul> <li>History of arthus-type hypersensitivity reactions after a previous dose of tetanus or diphtheria toxoid-containing vaccine; defer vaccination until at least 10 years have elapsed since the last tetanus-toxoid containing vaccine</li> </ul>
Varicella			Contraindications
			Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
			<ul> <li>Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, primary or acquired immunodeficiency, long-term immunosuppressive therapy, or patients with HIV infection who are severely immunocompromised)</li> </ul>
			Pregnancy
			Precautions
			Moderate or severe acute illness with or without fever
			Recent (within 11 months) receipt of antibody-containing blood product (specific interval depends on product)
			<ul> <li>Receipt of specific antivirals (i.e., acyclovir, famciclovir, or valacyclovir) 24 hours before vaccination; if possible, delay administration of these antiviral drugs for 14 days after vaccination.</li> </ul>
			Documented past history of Varicella disease.

IPV			Contraindications
			Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
			Precautions
			Pregnancy
			Moderate or severe acute illness with or without fever
Hepatitis B			Contraindications
			Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
			Precautions
			Infant weighing <2,000 grams if mother has documented hepatitis B surface antigen (HbsAg)-negative at the time of the infant's birth
			Moderate or severe acute illness with or without fever
MMR			Contraindications
			Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
			Pregnancy
			Known severe immunodeficiency (e.g., hematologic and solid tumors, or severely symptomatic infection)
			Precautions
			Recent (11 months) receipt of antibody-containing blood product (specific interval depends on product)
			History of thrombocytopenia or thrombocytopenic purpura
			Moderate or severe acute illness with or without fever
Meningococcal			Contraindications
			Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
			Precautions
			Moderate or severe acute illness with or without fever
			<ul> <li>Receipt of specific antivirals (acyclovir, famciclovir, or valacyclovir) 24 hours before vaccination; if possible, delay administration of these antiviral drugs for 14 days after vaccination.</li> </ul>
Haemophilus			Contraindications
influenza type b			Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
(Hib)			Age younger than 6 weeks
			Moderate or severe acute illness with or without fever
Pneumococcal			Contraindications
Disease			<ul> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose of PCV7 or PCV13 or to a vaccine component, including to any vaccine containing diphtheria toxoid</li> </ul>
			For PPSV23, severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component
			Precautions
			Moderate or severe acute illness with or without fever
imation of DPH,	my child has ndance at the	had, or is a childcare an	n of Public Health (DPH) declares an outbreak of a vaccine preventable disease, or if in the trisk of having an exposure to a vaccine preventable disease, my child shall be temporand/or school until the risk period ends, which may be three weeks or longer. My child shall by DPH.
Name of Pa	tient		DOB/
Name of Pa	rent/Guardian		
Signature (F	Patient/Parent	)	
ovider Informati		·	
			MD/DO/APRN/PA
Signature _			Date/
	Address		Phone #

School Vaccination Medical Exemption Form 073115